

**CUSD#303**  
**LIFE-THREATENING ALLERGY**  
**ACTION PLAN**

Student Name: \_\_\_\_\_ D.O.B: \_\_\_\_\_ Grade: \_\_\_\_\_

Allergy to: \_\_\_\_\_

Does this student have asthma?  Yes  No \*Higher risk for severe reaction

Place  
Child's  
Picture  
Here

**★ STEP 1: TREATMENT ★**

Symptoms	Give Checked Medication <small>(To be determined by physician authorizing treatment)</small>
• If exposure to allergen, but <i>no symptoms</i> :	<input type="checkbox"/> Epinephrine <input type="checkbox"/> Antihistamine
• <b>Mouth</b> Itching, tingling, or swelling of lips, tongue, mouth	<input type="checkbox"/> Epinephrine <input type="checkbox"/> Antihistamine
• <b>Skin</b> Hives, itchy rash, swelling of the face or extremities	<input type="checkbox"/> Epinephrine <input type="checkbox"/> Antihistamine
• <b>Gut</b> Nausea, abdominal cramps, vomiting, diarrhea	<input type="checkbox"/> Epinephrine <input type="checkbox"/> Antihistamine
• <b>Throat</b> †    Tightening of throat, hoarseness, hacking cough	<input type="checkbox"/> Epinephrine <input type="checkbox"/> Antihistamine
• <b>Lung</b> †      Shortness of breath, repetitive coughing, wheezing	<input type="checkbox"/> Epinephrine <input type="checkbox"/> Antihistamine
• <b>Heart</b> †     Weak or thready pulse, low blood pressure, fainting, pale, blueness	<input type="checkbox"/> Epinephrine <input type="checkbox"/> Antihistamine
• <b>Other</b> †	<input type="checkbox"/> Epinephrine <input type="checkbox"/> Antihistamine
• <b>If reaction is progressing in several of the above areas</b> <b>DO NOT HESITATE TO GIVE</b>	<input type="checkbox"/> Epinephrine <input type="checkbox"/> Antihistamine

†Potentially life-threatening. The severity of symptoms can quickly change.

**MEDICATION/DOSAGE**

Epinephrine (Brand & Dose): \_\_\_\_\_

Antihistamine (Brand & Dose): \_\_\_\_\_

Other (e.g., inhaler-bronchodilator if asthma): \_\_\_\_\_

**IMPORTANT: Asthma inhalers and/or antihistamines cannot be relied upon to replace epinephrine in anaphylaxis**

This student is authorized to self-carry/self-administer an Epinephrine auto-injector ?  Yes  No

**★ STEP 2: EMERGENCY CALLS ★**

1. Call 911. State that an allergic reaction has been treated, and additional epinephrine may be needed.

2. Licensed Health Care Provider: \_\_\_\_\_ Office Number: \_\_\_\_\_

3. Parents/Guardian: \_\_\_\_\_ Home: \_\_\_\_\_

Mother Cell: \_\_\_\_\_ Mother Work: \_\_\_\_\_

Father Cell: \_\_\_\_\_ Father Work: \_\_\_\_\_

**EVEN IF PARENT/GUARDIAN CANNOT BE REACHED, DO NOT HESITATE TO MEDICATE  
OR TAKE CHILD TO MEDICAL FACILITY!**

Parent/Guardian's Signature \_\_\_\_\_ Date \_\_\_\_\_

Health Care Provider Signature \_\_\_\_\_ Date \_\_\_\_\_

**\*\*COMPLETE BOTH SIDES\*\***

# Individual Life-Threatening Allergy Care Plan

## General Plan:

- Epinephrine will be stored in the:     Nurses Office     Class Room     On Person
- If applicable, the *Self-Carry/Self-Administer Waiver* form has been signed & submitted.
- Student's symptoms of an allergic reaction include: \_\_\_\_\_
- Student can recognize an allergic reaction and knows when and how to seek help.
- Classroom teachers will be made aware of plan.

## Bus Transportation:

- Drivers do not carry epinephrine or administer epinephrine.
- All drivers are trained to recognize symptoms of respiratory distress and will pull over and call 911 in the event of an emergency.

## Field Trip Procedures:

- Special needs will be identified prior to any off-campus trip.
- Prescribed medication & Emergency Action Plan must be reviewed and carried by certified staff member.

**Other Needs:** \_\_\_\_\_

## Parent/Guardian Plan:


- I give Health Services staff permission to communicate with the Health Care Provider about this medication.
- I understand that these medications may be administered by a certified staff member who has reviewed this care plan and the use of emergency medication.
- I agree that this information will be shared with school staff working with my child and 911 personnel, if needed.
- I assume responsibility for supplying medication that will not expire during the course of its intended use.
- I will provide medication in the original prescription container with instructions by above health care provider.
- If my child is authorized to self-carry, additional medication will be kept in the health office as recommended.

**Parent/Guardian Signature:** \_\_\_\_\_ **Date:** \_\_\_\_\_

**Reviewed by School Nurse:** \_\_\_\_\_ **Date:** \_\_\_\_\_

### EpiPen (Epinephrine) Auto-injector Directions

1. Remove the EpiPen Auto-injector from the plastic carrying case.
2. Pull off the blue safety release cap.
3. Swing and firmly push orange tip against mid-outer thigh.
4. Hold for approximately 10 seconds.
5. Remove and massage the area for 10 seconds.



### Auvi-Q (Epinephrine) Injection Directions

1. Remove the outer case of Auvi-Q. This will automatically activate voice instructions.
2. Pull off red safety guard.
3. Place black end against mid-outer thigh.
4. Press firmly and hold for 5 seconds.
5. Remove from thigh.

